## **Open Access**



# Low impact of *Zostera marina* meadows on sediment and water microbiota under brackish conditions

Daniel P. R. Herlemann<sup>1,4\*</sup>, Luis F. Delgado<sup>2</sup>, David J. Riedinger<sup>1</sup>, Víctor Fernández-Juárez<sup>3</sup>, Anders F. Andersson<sup>2</sup>, Christian Pansch<sup>5</sup>, Lasse Riemann<sup>3</sup>, Mia M. Bengtsson<sup>6</sup>, Greta Gyraitė<sup>7,8</sup>, Marija Kataržytė<sup>7</sup>, Veljo Kisand<sup>4</sup>, Sandra Kube<sup>1</sup>, Georg Martin<sup>9</sup>, Kasia Piwosz<sup>10</sup>, Marcin Rakowski<sup>10</sup> and Matthias Labrenz<sup>1</sup>

### Abstract

**Background** Zostera marina is an important ecosystem engineer influencing shallow water environments and possibly shaping the microbiota in surrounding sediments and water. Z. marina is typically found in marine systems, but it can also proliferate under brackish conditions. Changes in salinity generally have a strong impact on the biota, especially at the salty divide between salinity 6 and 9. To better understand the impact of the salty divide on the interaction between Z. marina and the surrounding sediment and water microbiota, we investigated the effects of Z. marina meadows on the surrounding microbiota across a salinity range of 6–15 in the Baltic Sea during the summer using 16S and 18S rRNA gene amplicon sequencing.

**Results** Salinity was the most important factor for structuring the microbiota within both water and sediment. The presence of *Z. marina* affected the composition of the bacterial and eukaryotic community and bacterial alpha diversity in the sediment. However, this effect was confined to alpha-mesohaline conditions (salinity 9–15). The impact of *Z. marina* below salinity 9 on water and sediment microbiota was insignificant.

**Conclusions** Increasing salinity was associated with a longer leaf length of *Z. marina*, causing an increased canopy height, which affects the sediment microbiota through reduced water velocity. Hence, we propose that the canopy effect may be the major predictor explaining *Z. marina*'s interactions with the surrounding microbiota at salinity 9–15. These findings emphasize the importance of the physical effects of *Z. marina* meadow ecosystem services and have important implications for *Z. marina* management under brackish conditions in a changing climate.

**Keywords** Coastal zone, Salinity, Horohalinicum, Baltic Sea, Bacterial community, Microeukaryotic community, Seagrass, Littoral, Eelgrass

\*Correspondence: Daniel P. R. Herlemann daniel.herlemann@io-warnemuende.de

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this articles are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

#### Introduction

In shallow, sheltered coastal regions, Zostera marina (also called eelgrass) builds large meadows that provide ecosystem services, including sediment stabilization, food provision, shelter, and nutrient recycling. It maintains intimate ecological interactions with microbial consortia that live in association with plants and within the surrounding seawater and sediments [1, 2]. Despite their ecological importance, Z. marina meadows decline globally by 2-5% annually, primarily due to human pressures, including climate change [3, 4]. Z. marina is widespread along coastlines throughout the Northern Hemisphere and has been suggested as a model system for aquatic plants [5]. Because the leaves and roots are constantly submerged, microorganisms attached to Z. marina interact strongly with the surrounding water and sediment bacterial communities [2, 6]. For example, the release of organic molecules (e.g., amino acids and sugars) and gases  $(CO_2, N_2)$  by aquatic macrophytes is associated with the presence of specific sulfate-reducing and nitrogen-fixing bacterial communities [7–10]. However, interactions with the surrounding environment extend beyond directly attached bacteria and may also be recognized within the Z. marina meadow, e.g., by the bacterial community in the sediment surrounding the roots [11-13] as well as the edges of *Z. marina* meadow patches [5].

In contrast to the effects on sediment, the influence of Z. marina on bacteria in the water columns of its meadows is still debated and seems to depend on environmental conditions [14–18]. For instance, based on a reduction in colony-forming bacterial units in the intertidal regions of tropical islands, Lamb et al. [19] suggested that Z. marina acts as a natural water filtration system for allochthonous pathogenic bacteria. Similarly, other studies have found lower levels of potentially pathogenic bacteria in Z. marina meadows than in areas without Z. marina [20–23]. Tasdemir et al. [24] and Millan et al. [25] reported the presence of potentially antibiotic-producing bacteria in Z. marina leaves, suggesting an influence on the surrounding seawater. However, yet other studies did not provide conclusive evidence of pathogen reduction by Z. marina in coastal waters [14, 15, 26]. Hence, the interaction between Z. marina and the surrounding bacterial community is not fully understood [14–16, 18].

In this study, we characterized the water and sediment microbiota inside, on the edge of, and outside of *Z. marina* meadows at different salinity levels at Baltic Sea coastal sites. *Z. marina* meadows are present within a salinity range of 5–35. Within this salinity range, especially at the salty divide at salinities 6–9, the sediment and water microbiota change from marine-related to freshwater-related organisms [27–31]. The salinity-driven shift in the biota at the salty divide [29, 32, 33] is a fundamental concept in ecology that has effects on

species richness and community composition in different biological groups [34]. Few studies have investigated the effects of the host and its surrounding microbiota among the salty divide [35, 36]. Previous studies investigating the impact of salinity on bacterial communities associated with macrophytes suggested that host and habitat were the most important factors in structuring their composition [37, 38], with the strongest changes occurring in the salty divide [39, 40]. We hypothesized that (1)the direct effect of Z. marina on the sediment and water microbiota differs depending on the salinity and that (2) the effect of Z. marina on the meadow water and sediment microbiota decreases with increasing distance from the Z. marina meadow. To address these hypotheses, we investigated Z. marina meadows in the Baltic Sea, where a stable salinity gradient with a characteristic sediment and water bacterial community exists [29, 30] and where Z. marina is highly abundant along the salinity gradient [41].

#### Materials and methods

Samples were collected as described by Riedinger et al. [26] in shallow coastal areas of the Baltic Sea (Fig. 1A-C). Each sampling station consisted of three subsamples taken within the *Z. marina* meadow ("inside"), approximately 15 m from the meadow ("edge") and at least 100 m from the *Z. marina* meadow ("outside") (Fig. 1D). Samples for DNA extraction from water and sediment were collected from each substation. For water, 100 mL syringes were used to sample ca. 5 cm from the *Z. marina* and 20 cm above the sediment surface. Sediment samples were collected using sterile plastic tubes by scraping the upper 1 cm at each station.

The *Z. marina* densities were counted in triplicate  $20 \times 20$  cm squares. Leaf length was measured for 30 plants in each meadow. All samples were immediately transferred to a 4 °C cooler and stored (maximally 8 h) until processing. Salinity, temperature, and depth were measured using a CTD48M (Sea & Sun Technology). The dissolved oxygen and pH were measured using a HQ40D portables 2-channel multimeter. Phosphate (PO<sub>4</sub><sup>3-</sup>), nitrate (NO<sub>3</sub><sup>-</sup>), nitrite (NO<sub>2</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>), and silicate (SiO<sub>2</sub>) concentrations were measured using a Seal Analytical QuAAtro automated continuous flow analyzer (SEAL Analytical Ltd. Nordestedt. Germany). Chlorophyll-*a* (Chl-*a*) was measured fluorometrically using a 10- AU-005-CE fluorometer (Turner San Jose, USA).

To determine bacterial and eukaryotic community composition, water was filtered through 0.2  $\mu$ m polyvinylidene fluoride membrane filters (Merck, Darmstadt, Germany) and shock frozen in liquid nitrogen. DNA was extracted after bead beating using a DNeasy PowerSoil Pro Kit Pro (Qiagen. Hilden. Germany) following the manufacturer's protocol after grinding the filters. For the

RV-14

#### A. Baltic Sea

#### **B. Beta-mesohaline samples**





C. Alpha-mesohaline samples



D. Sampling inside, on the edge and outside of the meadow



Fig. 1 Sampling location (A) alpha-mesohaline stations colored by salinity; (B) overview of the Baltic Sea sites where the subgraphs are marked (C) betamesohaline stations colored by salinity. (D) Sampling scheme for each station. Samples were collected inside the *Zostera marina* field, on the edge (15 m), and outside (100 m) of the *Z. marina* field

sediment samples, a subsample (300 mg) of the sediment was transferred into bead-beating tubes. The samples were placed on ice, sonicated twice for 7 min, and then subjected to bead-beating for 30 s at 4 m/s on a vortex adaptor. Subsequently, the manufacturer's instructions were followed, and the DNA yield was quantified using the PicoGreen assay (Thermo Fisher, Waltham. USA). The 16S rRNA genes were amplified using V3–V4 primers [29], and the 18S rRNA genes were amplified by targeting the eukaryotic V4 region [42]. The samples were sequenced on a MiSeq (Illumina Inc. San Diego. CA. US) for the 16S and 18S rRNA genes by SciLifeLab/NGI (Solna. Sweden). Amplicon sequencing data were processed using the workflow described by Riedinger et

al. [26] (https://github.com/biodiversitydata-se/amplico n-multi-cutadapt). ASVs with similar taxonomic assignment were aggregated.

For the bacterial community, sequences assigned to Archaea, chloroplasts, and mitochondria were removed before analysis because the primers used covered only minor parts of these groups. After the removal of these groups, a total of 11,599,548 reads were available for the bacterial community (Appendix Fig. 1A). For the eukaryotic data, reads assigned to unclassified Opisthokonta, Fungi, Embryophyceae (including *Zostera* spp.), and Metazoa were excluded, resulting in a total of 20,702,520 reads (Appendix Fig. 1B).

#### Statistical analysis

The amplicon sequence variant (ASV) richness ( $S_{OBS}$ ) and Shannon index were estimated using Explicet [43], which performs rarefaction-based analysis via bootstrapping. For all stations, bootstrap resampling was conducted at the size of the smallest library at the rarefaction point (Bacteria: 10,257 excluding two samples; Eukarya 25,417 excluding two samples) to compare ASVs between libraries with equal sampling efforts. The Kruskal–Wallis test and a *post hoc* Tukey's pairwise test was used to determine significant differences between the numbers of ASVs in the samples. A sequential Bonferroni correction was used for multiple comparisons. For the sample groups, p > 0.01 indicated that the variances were not significantly different from each other.

Due to the large spread between individual sampling sequencing depth (Bacteria: 8,720-138,054 reads, Eukarya: 9,699-213,379 reads), quality-trimmed sequencing reads were transformed using a centered log-ratio (CLR) transformation. This compositional data analysis approach [44] separates data variance according to differences rather than abundances. PCA on the pairwise Aitchison distances (Euclidean distance on CLR-transformed data [45]) was used as an exploratory analysis of the microbiota composition [46]. Then, the data were split into sub-datasets, and each micro-environment was tested separately to assess the impact of environmental factors. The effects of environmental variables on the community composition patterns of water and sediment were analyzed using PERMANOVA and two-way PERMANOVA based on Aitchison distances. The number of permutations was set to 9999, and p < 0.01was considered significant.

For analyzing differences between microbiota at different water depths, the averages of *Z. marina* abundance and length at the different sampling stations were calculated, with values above the average categorized as high and below the average as low (Appendix Table 1).

#### Results

#### Characterization of sampling sites

The samples were categorized following a modified Venice system [47], which takes the salty divide into account (Table 1): alpha-mesohaline conditions (salinity 9.3–15.5; stations BV5, BV6, BV7, BV8, BV10, BV22, BV24) and beta-mesohaline conditions (salinity 6.1-7.7; stations BV9, BV11, BV12, BV14, BV15, BV16, BV18, BV21). Both alpha- and beta-mesohaline sites were characterized by an average depth of ~2.6 m, low waves (~0.2 m), and wind speed (2.1 Bft). The average temperature and salinity were higher at alpha-mesohaline conditions compared to beta-mesohaline conditions. Chl-a, POC, PON, and SiO2 were, on average, more than twice as high under beta-mesohaline conditions than under alpha-mesohaline conditions. In particular, the beta-mesohaline station BV11 (Greifswalder Bodden) where higher compared to the average chemical measurements for Chl-a (19.7 mg m<sup>-3</sup>), PON (34.9 µmol L<sup>-1</sup>), POC (265.3 µg L<sup>-1</sup>), DOC (614.8  $\mu$ mol L<sup>-1</sup>) and SiO<sub>2</sub> (43.3  $\mu$ mol L<sup>-1</sup>) indicating the presence of a phytoplankton bloom.

# Microbial patterns in sediment and water within and outside the *Zostera marina* meadows

PCA of the bacterial (Fig. 2A) and eukaryotic (Fig. 2B) community composition showed a clear separation between the water and sediment microbiota along the first component (two-way PERMANOVA: Bacteria F = 131, p < 0.01; Eukarya F = 87, p < 0.01), independent of all other factors measured. Along the second component, a separation between alpha- and beta-mesohaline samples was apparent (two-way PERMANOVA: Bacteria F = 19, p < 0.01; Eukarya F = 16, p < 0.01). The interaction between the sediment/water and salinity category was also significant (two-way PERMANOVA: Bacteria F = 15, p < 0.01; Eukarya F = 11, p < 0.01). Because the sediment and water microbiota were clearly separated, we analyzed them separately. Envfit analysis of the sediment and water microbiota showed that the community composition was strongest associated (highest  $r^2$ ) with salinity (Table 2).

To examine the potential influence of the presence of *Z. marina* in water and sediment microbiota, samples were divided according to proximity to *Z. marina* meadows (inside, edge, and outside) and analyzed separately for alpha- and beta-mesohaline conditions and for water and sediment samples. PC analysis and PERMANOVA revealed no significant effect of proximity to *Z. marina* on the water bacterial and eukaryotic community composition under neither alpha (F=0.9, p=0.54) nor beta (F=1.1, p=0.23)-mesohaline conditions (Appendix Fig. 2C, D, G, H). The interaction effect between salinity and proximity to *Z. marina* was also nonsignificant (two-way PERMANOVA: F=0.59, p=0.72). Similarly, no significant differences in the water column alpha-diversity

nitrogen, NA=	not avail	able)	)			-	-			)		-	5		
	Sample	Salinity	Chla (mgm <sup>-3</sup> )	Depth <sup>1</sup> (m)	Wave (m)	Wind (Bft)	PO <sub>4</sub> (µmol I <sup>-1</sup> )	Length <sup>2</sup> (cm)	Temp (°C)	POC (µmol I <sup>-1</sup> )	DOC (µmol I <sup>-1</sup> )	NH <sub>4</sub> (µmol I <sup>-1</sup> )	PON (µmol I <sup>-1</sup> )	DN (µmol I <sup>-1</sup> )	SiO <sub>2</sub> (µmol I <sup>-1</sup> )
β – meso-	BV-21	6.7	1.6	3.3	0.3	m	0.5	39	16.4	21.4	378.1	1.3	3.9	18.4	15.8
haline	BV-18	6.1	2.3	2.6	0	0	NA	22	18.8	36.2	385.8	NA	5.2	18.3	9.0
	BV-16	6.7	1.5	3.2	0.1	1.8	0.2	42	17.8	21.6	354.8	1.2	3.6	19.7	10.0
	BV-15	6.8	2.4	2.4	0.2	2	0.1	28	19.0	31.5	372.8	0.6	4.6	19.3	9.8
	BV-14	6.3	3.1	4.5	NA	NA	0.1	38	18.7	61.4	426.1	NA	6.6	21.1	9.4
	BV-12	7.0	NA	2.0	0	0.3	NA	44	ΝA	NA	NA	NA	NA	NA	NA
	BV-11	7.6	19.7	2.2	0.5	4	0.5	53	20.0	265.3	614.8	0.8	34.9	28.0	43.3
	BV-9	7.7	2.3	2.9	0.9	4	0.3	44	20.4	37.5	373.9	NA	5.3	18.7	13.7
Average		6.9	4.4	2.5	0.1	2.3	0.3	39	18.9	61.9	407.6	1.0	8.4	20.3	15.2
a- meso-	BV-24	9.6	1.3	2.0	0.1	2	0.3	38	21.0	22.4	353.0	NA	3.7	18.3	9.5
haline	BV-22	11.7	NA	1.2	0.1	2.7	NA	74	ΝA	NA	NA	NA	NA	NA	NA
	BV-10	9.3	2.9	4.1	0.3	S	0.2	109	21.2	28.2	371.5	2.1	3.8	17.3	5.1
	BV-8	9.5	1.4	2.9	0.1	-	0.1	75	21.3	16.2	361.4	0.7	2.7	18.0	5.3
	BV-7	9.8	2.1	2.5	0.1	S	0.4	48	21.4	23.3	356.7	0.8	3.7	19.1	9.8
	BV-6	10.5	1.9	3.7	NA	NA	0.1	85	19.4	59.9	334.5	0.9	6.4	17.1	4.3
	BV-5	15.4	0.7	1.2	NA	NA	0.1	60	18.2	17.4	323.7	1.0	2.8	15.6	2.4
Average		10.9	1.6	2.5	0.1	2.3	0.2	70	20.0	26.7	344.7	1.3	3.7	17.3	5.9
Total Average		8.7	3.1	2.7	0.2	2.1	0.3	53	19.4	45.4	378.2	1.2	6.2	18.9	10.9
<sup>1</sup> water depth; $^2Z_{0}$	ostera marinu	a shoot len	gth												

**Table 1** Physicochemical parameters measured at the stations (see Fig. 1A-C). The substations "inside, edge, and outside" (see Fig. 1D) were averaged because the differences were minimal. *Zostera marina* shoot length was determined from inside samples. (ChI a=Chlorophyll a, DOC= dissolved organic carbon, PON= particular organic matter, DN= dissolved



Fig. 2 Principal component analysis of the (A) Bacterial community and (B) Eukaryotic community based on Aitchison distances in sediment (brown) and water (blue). Dots indicate samples taken under alpha-mesohaline conditions (salinity 9.3–16), and circles samples taken under beta-mesohaline conditions (salinity 6-7.8)

**Table 2** Results of the envfit analysis. PC1 and PC2 show the relationships of the environmental variables with the first and second PCA axes (based Aitchison distance). R<sup>2</sup> is the variation explained by the multiple regression model. Pr(>r) indicates the significance of the multiple regression. Significance codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05. (A) bacterial community water samples and (B) bacterial community sediment samples (C) eukaryotic community sediment samples

	PC1	PC2	r <sup>2</sup>	Pr(> <i>r</i> )
( <b>A</b> )				
Temperature	-0.54	-0.84	0.07	0.016*
Salinity	-0.99	0.05	0.69	0.001***
PON	0.94	0.35	0.18	0.001***
Chl-a	0.99	0.17	0.17	0.001***
POC	0.83	0.56	0.26	0.001***
( <b>B</b> )				
Temperature	0.98	-0.19	0.24	0.001***
Salinity	-0.99	0.01	0.67	0.001***
PON	-0.52	-0.86	0.09	0.005**
Chl-a	-0.51	-0.86	0.10	0.001***
POC	-0.47	-0.88	0.08	0.005**
( <b>C</b> )				
Temperature	-0.36	-0.93	0.15	0.001***
Salinity	-0.99	-0.10	0.54	0.001***
PON	0.64	-0.76	0.46	0.001***
Chl-a	0.64	-0.77	0.46	0.001***
POC	0.65	-0.75	0.45	0.001***
( <b>D</b> )				
Temperature	-0.83	0.56	0.26	0.001***
Salinity	-0.95	0.30	0.73	0.001***
PON	0.20	-0.98	0.06	0.041*
Chl-a	-0.19	-0.98	0.06	0.058
POC	-0.17	-0.98	0.06	0.049**

(neither S<sub>OBS</sub> nor Shannon index) in relation to *Z. marina* proximity were detected under either alpha-mesohaline or beta-mesohaline conditions (Kruskal-Wallis test: p > 0.01; Fig. 3, Appendix 3).

In contrast, both the bacterial and eukaryotic community compositions of the sediment samples were significantly different inside the Z. marina meadow compared to on the edge or outside the meadows for the alphamesohaline samples (F = 2.7, p < 0.01; Fig. 4, Appendix 2 B, F). However, in the lower saline beta-mesohaline samples, the effect of Z. marina proximity was not observed on the microbiota (F = 1.0 p = 0.3; Appendix 2 A, E). The interaction effect between salinity and proximity to Z. marina was insignificant (two-way PERMANOVA: F = 0.84, p = 0.08). Similar to the effects on the community composition, the presence of Z. marina affected the alpha-diversity (S<sub>OBS</sub> and Shannon index) of the bacterial sediment community, only at alpha-mesohaline conditions, where a significantly higher S<sub>OBS</sub> was observed inside the Z. marina meadow than at the edge and outside (Kruskal-Wallis test: p < 0.01; Fig. 3). Interestingly, no significant differences were found in the S<sub>OBS</sub> and Shannon index between the inside-edge-outside samples of the eukaryotes for sediment or for the water samples (Kruskal-Wallis test: p > 0.01, Fig. 3, Appendix 3).

*Z. marina* leaf length differed significantly between alpha and beta-mesohaline conditions (p < 0.01, Appendix 4). Additionally, there was a significant difference in the bacterial and eukaryotic communities among the different *Z. marina* length categories (Table 3). In addition, *Z. marina* abundance had a significant influence (p < 0.01) on the microbiota in contrast to water depth.



Fig. 3 Alpha diversity (S<sub>obs</sub>) of the (A) bacterial community and (B) eukaryotic community



Fig. 4 Effect of Zostera marina at alpha-mesohaline conditions on the sedimental (A) bacterial community and (B) sediment eukaryotic community

**Table 3** PERMANOVA of the bacterial (A) and eukaryotic (B) community compositions at water depth and of *Z. marina* abundance and leaf length categories. (A)

	water depth		Z. marina abu	ndance	Z. marina leaf length	
	F-value	<i>p</i> -value	F-value	<i>p</i> -value	F-value	<i>p</i> -value
( <b>A</b> )						
sediment	1.73	0.039	3.33	> 0.001	3.35	> 0.001
water	1.60	0.018	2.61	> 0.001	2.31	> 0.001
( <b>B</b> )						
sediment	1.43	0.060	2.54	> 0.001	2.98	> 0.001
water	1.48	0.043	2.54	> 0.001	3.34	> 0.001



Fig. 5 Bacterial phyla/classes (A) and eukaryotic phyla/classes (B) with abundances > 1% in the sediment and water at alpha-mesohaline and betamesohaline conditions



Fig. 6 Heatmap of the relative abundances of (A) bacterial and (B) eukaryotic amplicon sequence variants (ASVs) > 1% in sediment and water at alphamesohaline and beta-mesohaline conditions. (unclas. = unclassified)

#### Differences in taxonomic composition

The most abundant (>1%) phyla/classes of the bacterial communities in our study included Bacteroidetes (12-31%), Alphaproteobacteria (7-33%), and Gammaproteobacteria (10-22%; Fig. 5). The sediment samples contained higher relative abundances of Gammaproteobacteria (sediment 18% vs. 13% water), Planctomycetota (sediment 8% vs. 2% water), and Desulfobacterota (sediment 8% vs. 1% water). Alphaprotoeobacteria (sediment 7% vs. 27% water), Bacteroideta (sediment 17% vs. 24% water), Myxococcota (sediment 5% vs. >1% water), Chloroflexi (sediment 4% vs. >1% water), and Cyanobacteria (sediment 9% vs. 16% water) differed. Gammaproteobacteria (beta-mesohaline 15% vs. 21% alpha-mesohaline), Actinobacteria (beta-mesohaline 19% vs. 12% alphamesohaline) and Bacteroideta (beta-mesohaline 14% vs. 20% alpha-mesohaline) differed between the alpha and beta-mesohaline conditions. Within the Z. marina

meadow under alpha-mesohaline conditions, a high abundance of Campylobacterota (4% inside vs. > 1% outside) was observed. The alpha-mesohaline and betamesohaline water bacterial community compositions also differed. Alphaproteobacteria (beta-mesohaline 21% vs. 32% alpha-mesohaline) and Bacteroidota (beta-mesohaline 18% vs. 30% alpha-mesohaline) were more abundant under alpha-mesohaline conditions, whereas Actinobacteria (beta-mesohaline 21% vs. 10% alpha-mesohaline) and Cyanobacteria (beta-mesohaline 22% vs. 9% alphamesohaline) were more abundant under beta-mesohaline conditions.

On a finer taxonomic level, *Cyanobium* was most abundant ASV in the water column (max = 21%) especially under beta-mesohaline conditions, but it was also abundant in all other habitats (Fig. 6A). The abundant bacterial groups typically found in the water column were SAR11 clades, NS3a, NS5, SAR86, and *Planktomarina* 

under alpha-mesohaline conditions, whereas hgcI (betamesohaline 9.3% vs. 2.1% alpha-mesohaline) and CL500-29 (beta-mesohaline 4.8% vs. 1.1% alpha-mesohaline) were more abundant under beta-mesohaline conditions. The bacterial communities in the sediments were dominated by *Cyanobium, Illumatobacter*, unclassified Saprospiraceae, and the desulfosarcinal Sva0081 group. The bacterial genus Campylobacterota with higher abundances inside the *Z. marina* meadow at alpha-mesohaline conditions was mostly assigned to *Sulfurovorum*. Despite having a low overall abundance (0.8%), the genus has an abundance of 4% within the meadow. Representatives of the genus *Sulfurovorum* are able to oxidize sulfur.

Similar to the bacterial community composition, the community of microbial eukaryotes differed among the sediment and water samples (Fig. 6B). In water, *Picochlorum* and *Monodus* were abundant genera at beta-meso-haline conditions. *Rhizosolenia delicatula* was abundant under alpha-mesohaline conditions in water and sediment (beta-mesohaline 0% vs. 4.7% and 13.2% alpha-mesohaline). *Choricystis* sp. were more abundant inside (inside 13.3% vs. 3.6% outside) the *Z. marina* meadow and Gymnodiniaceae, rather than outside the meadow (inside 0.3% vs. 12.0% outside).

#### Discussion

Z. marina meadows stabilize sediments, provide food and shelter, and recycle nutrients, providing critical aquatic ecosystem services. Z. marina has been suggested to impact its environment beyond the direct Z. marina leaves (phyllosphere) [14–17, 19] and roots (rhizosphere) [11–13]. Our results showed a limited effect of Z. marina on the surrounding water and sediment microbiota since only at alpha-mesohaline conditions in sediments a significant effect of Z. marina on the microbiota was observed. Therefore, the first hypothesis was partially supported. Previous studies also observed an influence of Z. marina under saltwater conditions [12– 14, 16, 17, 48–50].

The missing effect of *Z. marina* meadows at lower salinity could be related to the symbiont (microbiota) or host (*Z. marina*). Among the mechanisms by which *Z. marina* influences the surrounding microbiome is the release of bioavailable carbon and the resulting accumulation of microbial grazers [51] and bacteria that excrete antimicrobial substances [52]. However, these effects appear to be local and may be detected only when the phyllosphere or the rhizosphere is directly sampled. The release of oxygen from the roots of *Z. marina* has also been shown to influence the surrounding microbiota [6]. However, since we sampled oxygenated top-layer sediments, the impact of released oxygen was likely minor. Another proposed mechanism is the absorption of pathogens by biofilms [53], which was not observed in in lower salinities [26].

Mechanisms of Z. marina influencing the surrounding sediment and water include a reduction of wave energy and increase sedimentation as a physical effect of Z. marina canopy structure ("canopy effect") [54, 55]. Under beta-mesohaline conditions, Z. marina has a lower productivity that is connected with a significant reduced leaf length (Table 1, Appendix 4). Shorter leaves also reduce the canopy effect, which reduces the capacity of Z. marina to trap fine sediments, including particle-attached bacterial communities [56] from the water column. At higher salinities, longer leaves are more likely to trap particle-attached bacterial communities, resulting in a significant change in the meadow surface sediment bacterial community [18]. Accordingly, a correlation between the canopy structure of Z. marina and the microbiota was previously reported [5, 18, 55, 57]. However, other local factors, in addition to salinity, including sediment type, history, and wave exposure, can also influence leaf length [58, 59].

In contrast to previous studies, no specific bacterial community was found on the edge of a Z. marina meadow in the water and sediment [5, 18]. Therefore, the second hypothesis was rejected. This could be attributed to the fact that the edges of the Z. marina meadow in our study were predominantly patchy and sampled in a distance of 15 m from the main meadow. Samples closer to the meadow may show a different result. We also found higher sediment bacterial diversity within the Z. marina meadows than outside under alpha-mesohaline conditions. This increased diversity of microbiota in close proximity to plant roots due to exudates was previously described as the "rhizosphere effect" [60]. Other studies [5, 61, 62] in contrast suggested higher diversity in bulk sediment outside the Z. marina meadow because the rhizosphere selects microorganisms from the bare sediment and reduces overall microbial diversity. The discrepancy may be connected to the brackish conditions in our study compared with the saltwater conditions in other studies and the fact that we did not sample the rhizosphere, but rather surface sediment in the Z. marina meadow.

The missing "*Z. marina* meadow"-effect of the water bacterial and microeukaryotic communities is in line with studies on the San Diego coastal water, in which the location of the sampling site was identified as the major driver of the bacterial community composition rather than proximity to the *Z. marina* meadow [15]. Other studies have reported a reduction in colony-forming units in *Z. marina* beds compared with surrounding water [20–23]. However, these studies were conducted at higher salinity. The reduced canopy effect in our study may also have reduced the effects on the water bacterial community. In addition to the canopy effect, a potential mechanism by which *Z. marina* directly influences water microbial communities is the release of exudates as dissolved organic matter (DOM) [6, 63]. The high DOM concentration in the Baltic Sea [64], which is caused by terrestrial influences, especially in the littoral zone, may mask the effects of DOM release on aquatic microbial communities. Moreover, DOM can be confounded by salinity; therefore, more studies, including experiments under controlled conditions, are needed to support the observed trends in this study.

#### Other effects on the microbiota

The bacterial and eukaryotic communities in the sediment and water were significantly different (Fig. 2). This is in accordance with previous studies in which the effect of microenvironment type has been shown to have a stronger impact on determining the bacterial community composition than salinity [14, 17, 65-67]. The second most important factor influencing microbiota structure was salinity. Based on the salinity range covered (salinity 6-15), clustering of the samples according to alpha-mesohaline and beta-mesohaline conditions was expected [28-30, 34, 68]. The S<sub>OBS</sub>, which reflects microbial richness, was highest in the sediment (Fig. 3), which is consistent with previous studies [5, 29, 30]. In contrast to the species minimum at the salty divide ("horohalinicum") observed for macroorganisms [34] and phytoplankton [68], the bacterial and microeukaryotic community species richness in the water was rather constant at different salinities (Fig. 3). No significant effects on bacterial and microeukaryotic  $\mathbf{S}_{\text{OBS}}$  have been reported in previous studies [29, 31, 69, 70]. In accordance with these studies, a shift was observed between the dominance of clades present in saltwater, including SAR11 clade I, NS3a, NS5, SAR86, and Planktomarina, and the dominance of clades typically found in freshwater, including HGC-I, SAR11-III, and CL500-29, at the salty divide. Most ASVs were assigned only to their taxonomic genus due to the short-read length of the method and the lack of species descriptions in current databases [71]. This resulted in a relatively low diversity of taxonomic genera and many undescribed taxonomic clades (Fig. 6).

For the water microbiota, a rather gradual change in community composition was observed with changing salinity, whereas the distinction between the alpha- and beta-mesohaline microbial communities in sediments was more pronounced (Fig. 2). This could be connected with a higher resistance of sediment bacterial communities to changes in salinity [66] and stronger mixing of water masses than in sediment, but also with differences in the sediment characteristics of the two salinity conditions.

#### Conclusion

A global effort is underway to restore coastal and estuarine ecosystems, and a major part of this effort is the reestablishment of Z. marina [72]. Our findings provide fundamental information about the impact of Z. marina on sediment and water microbiota under brackish conditions. We found that the influence of Z. marina on the microbiota was lower under brackish conditions, especially below a salinity of 9. The shift in salinity has a significant effect on both the microbiota and host appearance. Because leaf length can affect the impact on the sediment bacterial community in a Z. marina meadow, we suggest that the canopy effect is pivotal to the influence of Z. marina meadows on the surrounding environment. This finding supports the previous hypothesis that the effects of Z. marina are rather passive [18, 55]. The results should be considered when managing brackish Z. marina meadow ecosystem services in brackish environments, such as the Baltic Sea, Black Sea, and larger estuaries.

#### Supplementary Information

The online version contains supplementary material available at https://doi.or g/10.1186/s40793-024-00662-6.

Supplementary Material 1

#### Acknowledgements

The authors acknowledge support from the National Genomics Infrastructure in Genomics Application Stockholm funded by the Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council, and SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure. We are grateful to the crew and captain of the R/V Elisabeth Mann Borgese (EMB283), the research divers of the Leibniz Institute for Baltic Sea Research, the University of Rostock, Klaipeda University, and the Estonian Marine Institute, to Jonas Nilsson for assisting with sampling in Sweden, and to Adam Woźniczka and Jarone Pinhassi for providing lab space. In addition, the Leibniz Institute for Baltic Sea Research Warnemünde is conducting research in the new research field "Shallow Water Processes and Transitions to the Baltic Scale."

#### Author contributions

Conceptualization by DH, AA, CP, LR, MK, SK, MR, MB, ML; Data Curation by all authors; Formal Analysis by DH, LD, DR, VJ, KK, AA, CP, LR, MK, SK, ML; Funding acquisition by DH, AA, CP, LR, MK, TR, MR, ML. Project administration by SK, ML; Writing original draft by DH, DR, ML; Writing, review, and edition by all authors; Investigation by DH, DR, KK, AA, CP, LR, MB, GG, SK, GM, MR, ML; Visualization by DH; Software by AA, LD, VK, DH; Validation by SK.

#### Funding

Open Access funding enabled and organized by Projekt DEAL. This work resulted from the BiodivERsA project "Pathogenic *Vibrio* bacteria in the current and future Baltic Sea waters: mitigating the problem" (BaltVib), funded by the European Union and the Estonian Research Council (grant T210076PKKH / P200028PKKH), the Federal Ministry of Education and Research, Germany (grant 16LC2022A), the Innovation Fund Denmark (grant 0156-00001B), the Research Council for Environment, Agricultural Sciences and Spatial Planning FORMAS (grant 2020–02366), the Polish National Science Center (grant 2020/02/V/NZ8/00009), and the Research Council of Finland (Academy of Finland; grant 344743). In addition, the Leibniz Institute for Baltic Sea Research

Warnemünde is conducting research in the new research field "Shallow Water Processes and Transitions to the Baltic Scale."

#### Data availability

The datasets supporting the conclusions of this article are available in the European Nucleotide Archive under accession number PRJEB68222, which is in compliance with the Minimum Information about any (X) Sequence (MIxS) standard through the brokerage service GFBio93. Environmental data are available from IOWMeta (doi.io-warnemuende.de/10.12754/data-2023-0010).

#### Declarations

**Ethics approval and consent to participate** Not applicable.

#### **Consent for publication**

All authors have approved the final manuscript.

#### **Competing interests**

The authors declare no competing interests.

#### Author details

<sup>1</sup>Biological Oceanography, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), 18119 Rostock, Germany

<sup>2</sup>Science for Life Laboratory, School of Biotechnology, Division of Gene Technology, KTH Royal Institute of Technology, Solna 171 21, Sweden <sup>3</sup>Department of Biology, University of Copenhagen, Helsingør 3000, Denmark

<sup>4</sup>Center for Limnology, Estonian University of Life Sciences, Tartu 51006, Estonia

<sup>5</sup>Faculty of Science and Engineering, Environmental and Marine Biology, Åbo Akademi University, Turku/Åbo 20500, Finland

<sup>6</sup>Institute of Microbiology, University of Greifswald, 17489 Greifswald, Germany

<sup>7</sup>Marine Research Institute, Klaipėda University, Klaipėda 92294, Lithuania <sup>8</sup>Institute of Biosciences, Life Sciences Center, Vilnius University, Vilnius 10257. Lithuania

<sup>9</sup>Estonian Marine Institute, University of Tartu, Tallinn 12618, Estonia <sup>10</sup>National Marine Fisheries Research Institute, Gdynia 81-332, Poland

#### Received: 26 July 2024 / Accepted: 24 December 2024 Published online: 11 January 2025

#### References

- Walters K. The effects of complex trophic interactions on a marine microbenthic community. Ecology. 1993;74(5):1475–89.
- Duarte CM, Holmer M, Marbà N. Plant–microbe interactions in seagrass meadows. Interact between macro-and Microorganisms Mar Sediments. 2005;60:31–60.
- Dunic JC, Brown CJ, Connolly RM, Turschwell MP, Côté IM. Long-term declines and recovery of meadow area across the world's seagrass bioregions. Glob Change Biol. 2021;27(17):4096–109.
- 4. Duarte CM. The future of seagrass meadows. Environ Conserv. 2002;29(2):192–206.
- Ettinger CL, Voerman SE, Lang JM, Stachowicz JJ, Eisen JA. Microbial communities in sediment from *Zostera marina* patches, but not the Z. marina leaf or root microbiomes, vary in relation to distance from patch edge. PeerJ. 2017;5:e3246.
- Ugarelli K, Chakrabarti S, Laas P, Stingl U. The seagrass holobiont and its microbiome. Microorganisms. 2017;5(4):81.
- Donnelly A, Herbert R. Bacterial interactions in the rhizosphere of seagrass communities in shallow coastal lagoons. J Appl Microbiol. 1998;85(S1):S151–60.
- Nielsen LB, Finster K, Welsh DT, Donelly A, Herbert RA, De Wit R, Lomstein BA. Sulphate reduction and nitrogen fixation rates associated with roots, rhizomes and sediments from Zostera noltii and Spartina maritima meadows. Environ Microbiol. 2001;3(1):63–71.

- Pollard PC, Moriarty D. Organic carbon decomposition, primary and bacterial productivity, and sulphate reduction, in tropical seagrass beds of the Gulf of Carpentaria, Australia. Mar Ecol Progress Ser Oldendorf. 1991;69(1):149–59.
- Liu P, Zou S, Zhang H, Liu Q, Song Z, Huang Y, Hu X. Genome-resolved metagenomics provides insights into the microbialmediated sulfur and nitrogen cycling in temperate seagrass meadows. Front Mar Sci. 2023;10:1245288.
- Sun Y, Song Z, Zhang H, Liu P, Hu X. Seagrass vegetation affect the vertical organization of microbial communities in sediment. Mar Environ Res. 2020;162:105174.
- 12. Zheng P, Wang C, Zhang X, Gong J. Community structure and abundance of archaea in a Zostera marina meadow: a comparison between seagrass-colonized and bare sediment sites. Archaea. 2019;2019.
- Jensen SI, K
  ühl M, Prieme A. Different bacterial communities associated with the roots and bulk sediment of the seagrass Zostera marina. FEMS Microbiol Ecol. 2007;62(1):108–17.
- 14. Crump BC, Koch EW. Attached bacterial populations shared by four species of aquatic angiosperms. Appl Environ Microbiol. 2008;74(19):5948–57.
- Webb SJ, Rabsatt T, Erazo N, Bowman JS. Impacts of Zostera eelgrasses on microbial community structure in San Diego coastal waters. Elementa: Science of the Anthropocene. 2019;7.
- Crump BC, Wojahn JM, Tomas F, Mueller RS. Metatranscriptomics and amplicon sequencing reveal mutualisms in seagrass microbiomes. Front Microbiol. 2018;9:388.
- Fahimipour AK, Kardish MR, Lang JM, Green JL, Eisen JA, Stachowicz JJ. Global-scale structure of the eelgrass microbiome. Appl Environ Microbiol. 2017;83(12):e03391–16.
- Iqbal MM, Nishimura M, Sano M, Yoshizawa S. Particle-attached microbes in Eelgrass Vegetation areas Differ in Community structure depending on the Distance from the Eelgrass Bed. Microbes Environ. 2023;38(3):ME23013.
- Lamb JB, Van De Water JA, Bourne DG, Altier C, Hein MY, Fiorenza EA, et al. Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes, and invertebrates. Science. 2017;355(6326):731–3.
- Palazón A, Aragonés L, López I, López-Úbeda R, Saval Pérez JM. Determination of the most influential factors in the concentration of bacteria in coastal waters. 2018.
- 21. Deng Y, Liu S, Feng J, Wu Y, Mao C. What drives putative bacterial pathogens removal within seagrass meadows? Mar Pollut Bull. 2021;166:112229.
- Sun H, Wang T, Liu S, Tang X, Sun J, Liu X, et al. Novel insights into the rhizosphere and seawater microbiome of *Zostera marina* in diverse mariculture zones. Microbiome. 2024;12(1):27.
- Reusch TB, Schubert PR, Marten S-M, Gill D, Karez R, Busch K, Hentschel U. Lower Vibrio spp. abundances in Zostera marina leaf canopies suggest a novel ecosystem function for temperate seagrass beds. Mar Biol. 2021;168:1–6.
- 24. Tasdemir D, Scarpato S, Utermann-Thüsing C, Jensen T, Blümel M, Wenzel-Storjohann A, et al. Epiphytic and endophytic microbiome of the seagrass *Zostera marina*: do they contribute to pathogen reduction in seawater? Sci Total Environ. 2024;908:168422.
- 25. Millan C. The effect of seagrass components on the associated bacterial diversity and antimicrobial activity in *Zostera marina*. 2022.
- Riedinger DJ, Fernández-Juárez V, Delgado LF, Sperlea T, Hassenrück C, Herlemann DP, et al. Control of Vibrio vulnificus proliferation in the Baltic Sea through eutrophication and algal bloom management. Commun Earth Environ. 2024;5(1):246.
- Crump BC, Armbrust EV, Baross JA. Phylogenetic analysis of particle-attached and free-living bacterial communities in the Columbia River, its estuary, and the adjacent coastal ocean. Appl Environ Microbiol. 1999;65(7):3192–204.
- 28. Lozupone CA, Knight R. Global patterns in bacterial diversity. Proceedings of the National Academy of Sciences. 2007;104(27):11436-40.
- Herlemann DP, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. ISME J. 2011;5(10):1571–9.
- Klier J, Dellwig O, Leipe T, Jürgens K, Herlemann DP. Benthic bacterial community composition in the oligohaline-marine transition of surface sediments in the Baltic Sea based on rRNA analysis. Front Microbiol. 2018;9:236.
- Mazur-Marzec H, Andersson AF, Błaszczyk A, Dąbek P, Górecka E, Grabski M, et al. Biodiversity of microorganisms in the Baltic Sea: the power of novel methods in the identification of marine microbes. FEMS Microbiol Rev. 2024;48(5):fuae024.
- Paver SF, Muratore D, Newton RJ, Coleman ML. Reevaluating the salty divide: phylogenetic specificity of transitions between marine and freshwater systems. Msystems 2018;3(6). https://doi.org/10.1128/msystems.00232-18.
- 33. Kinne O. Salinity-invertebrates. Marine ecology. 1971.

- Remane A. Die Brackwasserfauna. Verhandlungen Der Deutschen Zoologischen Gesellschaft. 1934;36:34–74.
- Kivistik C, Käiro K, Tammert H, Sokolova IM, Kisand V, Herlemann DP. Distinct stages of the intestinal bacterial community of *Ampullaceana Balthica* after salinization. Front Microbiol. 2022;13:767334.
- Kivistik C, Knobloch J, Käiro K, Tammert H, Kisand V, Hildebrandt J-P, Herlemann DP. Impact of salinity on the gastrointestinal bacterial community of *Theodoxus fluviatilis*. Front Microbiol. 2020;11:683.
- Hempel M, Blume M, Blindow I, Gross EM. Epiphytic bacterial community composition on two common submerged macrophytes in brackish water and freshwater. BMC Microbiol. 2008;8:1–10.
- Behera P, Mahapatra S, Mohapatra M, Kim JY, Adhya TK, Raina V, et al. Salinity and macrophyte drive the biogeography of the sedimentary bacterial communities in a brackish water tropical coastal lagoon. Sci Total Environ. 2017;595:472–85.
- Van der Loos LM, D'hondt S, Engelen AH, Pavia H, Toth GB, Willems A, et al. Salinity and host drive Ulva-associated bacterial communities across the Atlantic–Baltic Sea gradient. Mol Ecol. 2023;32(23):6260–77.
- Mohapatra M, Manu S, Dash SP, Rastogi G. Seagrasses and local environment control the bacterial community structure and carbon substrate utilization in brackish sediments. J Environ Manage. 2022;314:115013.
- Asmala E, Gustafsson C, Krause-Jensen D, Norkko A, Reader H, Staehr PA, Carstensen J. Role of eelgrass in the coastal filter of contrasting Baltic Sea environments. Estuaries Coasts. 2019;42:1882–95.
- Hugerth LW, Muller EE, Hu YO, Lebrun LA, Roume H, Lundin D, et al. Systematic design of 18S rRNA gene primers for determining eukaryotic diversity in microbial consortia. PLoS ONE. 2014;9(4):e95567.
- Robertson CE, Harris JK, Wagner BD, Granger D, Browne K, Tatem B, et al. Explicet: graphical user interface software for metadata-driven management, analysis and visualization of microbiome data. Bioinformatics. 2013;29(23):3100–1.
- 44. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome datasets are compositional: and this is not optional. Front Microbiol. 2017;8:2224.
- Aitchison J. Principal component analysis of compositional data. Biometrika. 1983;70(1):57–65.
- Ramette A. Multivariate analyses in microbial ecology. FEMS Microbiol Ecol. 2007;62(2):142–60. https://doi.org/10.1111/j.1574-6941.2007.00375.x
- den Hartog C. Typologie Des Brackwassers. Helgoländer Wissenschaftliche Meeresuntersuchungen. 1964;10(1):377–90.
- Cúcio C, Engelen AH, Costa R, Muyzer G. Rhizosphere microbiomes of European seagrasses are selected by the plant, but are not species specific. Front Microbiol. 2016;7:440.
- 49. Sun F, Zhang X, Zhang Q, Liu F, Zhang J, Gong J. Seagrass (Zostera marina) colonization promotes the accumulation of diazotrophic bacteria and alters the relative abundances of specific bacterial lineages involved in benthic carbon and sulfur cycling. Appl Environ Microbiol. 2015;81(19):6901–14.
- Markovski M, Najdek-Dragić M, Herndl GJ, Korlević M. Compositional stability of sediment microbial communities during a seagrass meadow decline. 2022.
- Vymazal J. Removal of enteric bacteria in constructed treatment wetlands with emergent macrophytes: a review. J Environ Sci Health. 2005;40(6–7):1355–67.
- 52. Sundaravadivel M, Vigneswaran S. Constructed wetlands for wastewater treatment. Crit Rev Environ Sci Technol. 2001;31(4):351–409.
- Stott R, Tanner C. Influence of biofilm on removal of surrogate faecal microbes in a constructed wetland and maturation pond. Water Sci Technol. 2005;51(9):315–22.
- 54. Agawin NS, Duarte CM. Evidence of direct particle trapping by a tropical seagrass meadow. Estuaries. 2002;25:1205–9.

- Liu S, Wu Y, Luo H, Ren Y, Jiang Z, Zhang X, et al. Seagrass canopy structure mediates putative bacterial pathogen removal potential. Front Mar Sci. 2023;9:1076097.
- Rieck A, Herlemann DP, Jürgens K, Grossart H-P. Particle-associated differ from free-living bacteria in surface waters of the Baltic Sea. Front Microbiol. 2015;6:1297.
- Iqbal MM, Nishimura M, Haider MN, Yoshizawa S. Microbial communities on eelgrass (Zostera marina) thriving in Tokyo Bay and the possible source of leaf-attached microbes. Front Microbiol. 2023;13:1102013.
- Stevenson A, O Corcora TC, Hukriede W, Schubert PR, Reusch TB. Substantial seagrass blue carbon pools in the southwestern Baltic Sea include relics of terrestrial peatlands. Front Mar Sci. 2022;9:Art. Nr. 949101.
- 59. Salo T, Pedersen MF, Boström C. Population specific salinity tolerance in eelgrass (Zostera marina). J Exp Mar Biol Ecol. 2014;461:425–9.
- Dotaniya M, Meena V. Rhizosphere effect on nutrient availability in soil and its uptake by plants: a review. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences. 2015;85:1–12.
- 61. Ling N, Wang T, Kuzyakov Y. Rhizosphere bacteriome structure and functions. Nat Commun. 2022;13(1):836.
- Leach JE, Triplett LR, Argueso CT, Trivedi P. Communication in the phytobiome. Cell. 2017;169(4):587–96.
- Chen PW-Y, Annabel CN, Olivia M, Chou W-C, Chen J-J, Mukhanov V et al. Investigation of the Growth and Mortality of Bacteria and Synechococcus spp. in Unvegetated and Seagrass Habitats. 2024.
- Seidel M, Manecki M, Herlemann DP, Deutsch B, Schulz-Bull D, Jürgens K, Dittmar T. Composition and transformation of dissolved organic matter in the Baltic Sea. Front Earth Sci. 2017;5:31.
- Han Z, Wang C, Lei B, Hui N, Yu Y, Shi Y, Zheng J. A limited overlap of interactions between the bacterial community of water and sediment in wetland ecosystem of the Yellow River floodplain. Front Microbiol. 2023;14:1193940.
- Tammert H, Kivistik C, Kisand V, Käiro K, Herlemann DP. Resistance of freshwater sediment bacterial communities to salinity disturbance and implication for industrial salt discharge and climate change based salinization. Front Microbiomes. 2024;2:1232571.
- 67. Gordon-Bradley N, Lymperopoulou DS, Williams HN. Differences in bacterial community structure on Hydrilla verticillata and Vallisneria americana in a freshwater spring. Microbes Environ. 2014;29(1):67–73.
- Olli K, Ptacnik R, Klais R, Tamminen T. Phytoplankton species richness along coastal and estuarine salinity continua. Am Nat. 2019;194(2):E41–51.
- Herlemann DP, Lundin D, Andersson AF, Labrenz M, Jürgens K. Phylogenetic signals of salinity and season in bacterial community composition across the salinity gradient of the Baltic Sea. Front Microbiol. 2016;7:1883.
- Telesh IV, Schubert H, Skarlato SO. Revisiting Remane's concept: evidence for high plankton diversity and a protistan species maximum in the horohalinicum of the Baltic Sea. Mar Ecol Prog Ser. 2011;421:1–11.
- Glöckner FO, Yilmaz P, Quast C, Gerken J, Beccati A, Ciuprina A, et al. 25 years of serving the community with ribosomal RNA gene reference databases and tools. J Biotechnol. 2017;261:169–76.
- Unsworth RK, Jones BL, Coals L, Furness E, Inman I, Rees SC, Evans AJ. Overcoming ecological feedbacks in seagrass restoration. Restor Ecol. 2024;32(4):e14101.

#### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.